



Class: FY BSc

Subject : Calculus

Subject Code: PUSASQF1.1

Chapter: Unit 2 Chapter 1

Chapter Name: Derivatives

Today's Agenda

1. Derivatives
 1. Derivatives Notation
 2. Why to study derivatives?
 3. Application of Derivatives in daily life
2. Differentiation Formulas
3. Higher Order Derivatives
4. Differentiation Rules
5. Chain Rule of Differentiation
6. Partial Differentiation

1 Derivative



- Derivative, in mathematics is defined as the rate of change of a function with respect to a variable.
- The derivative of a function of a single variable at a chosen input value, when it exists, is the slope of the tangent line to the graph of the function at that point.
- The derivative is often described as the "instantaneous rate of change", the ratio of the instantaneous change in the dependent variable to that of the independent variable.

Let $f(x)$ be a function defined in an open interval containing a . The derivative of the function $f(x)$ at a , denoted by $f'(a)$, is defined by :

$$f'(a) = \lim_{x \rightarrow a} \left(\frac{f(x) - f(a)}{x - a} \right)$$



<https://www.youtube.com/watch?v=rAof9Ld5sOg>

1.1 Derivatives notation

Derivatives are the result of performing a differentiation process upon a function or an expression. Derivative notation is the way we express derivatives mathematically.

1. Lagrange's notation :

In Lagrange's notation, the derivative of f is expressed as f' prime (pronounced "*f prime*"). This notation is probably the most common when dealing with functions with a single variable.

2. Leibniz's notation

In Leibniz's notation, the derivative of f is expressed as $\frac{d}{d(x)} f(x)$

This notation, becomes very useful when dealing with integral calculus, differential equations, and multivariable calculus.

3. Newton's notation

In Newton's notation, the derivative of f is expressed as $\dot{f}$ and the derivative of $y = f(x)$ is expressed as $\dot{y}$. This notation is mostly common in Physics and other sciences where calculus is applied in a real-world context.

1.2 Why to study derivatives ?

- The original purpose of the derivative is to analyze the sensitivity or rate of change of a function with respect to its independent variable — that is, given a tiny change in the independent variable x , how much does the dependent variable, y , respond to that change. If the derivative of a function is high, it means that, for that function, the dependent variable responds greatly to that tiny change in its independent variable; if the derivative is low, the dependent variable changes a very small amount in response.
- When taking the derivative of a function, you can find relative extrema, critical values, and where the function is increasing decreasing. Taking the derivative again, you can find the concavity of the function, and the points of inflection that exist on it.
- Derivatives are used in velocity and acceleration in physics, marginal profit functions in business, and growth rates in biology.

1.3 Application of derivatives in daily life

The derivative can be used for many other purposes, such as **analyzing the increasing/decreasing behavior and concavity of a function.**

Example:

Let's say the population in an island at a certain time t can be modeled by the function $P(t)$. If, say, the population is increasing, we know that after a tiny change in time can result in a tiny, positive change in population; if the population is decreasing, a tiny change in time results in a tiny, negative change in population.

Using what we discussed above, the derivative of P , can tell us the increasing / decreasing behavior of population with its sign.

If $\frac{dP}{dt}$ is positive, it means that a tiny change in time is accompanied by a tiny positive growth in

population; the same logic goes with negative.

Hence, if the derivative is positive, the original function is increasing; if the derivative is negative, the original function is decreasing.

2 Differentiation Formulas

1. $\frac{dk}{dx} = 0$ where $k = \text{constant}$

2. $\frac{d(x)}{dx} = 1$

3. $\frac{d(kx)}{dx} = k$ where $k = \text{constant}$

2 Example

Example 1:

$$\frac{d}{dx}(6) = 0$$

Example 2:

$$\frac{d}{dx}(5x + 4) = 5(1) + 0 = 5$$

2 Differentiation Formulas

$$4. \frac{d(x^n)}{dx} = nx^{n-1}$$

$$5. \frac{d}{dx} \sqrt{x} = \frac{1}{2\sqrt{x}}$$

2 Example

Example 1:

$$\frac{d}{dx} x^8 = 8x^7$$

Example 2:

$$\frac{d}{dx} \frac{6}{x^4} = 6(-4) x^{-4-1} = -24 x^{-5}$$

2 Differentiation Formulas

$$6. \frac{d(e^x)}{dx} = e^x$$

$$7. \frac{d(\ln(x))}{dx} = \frac{1}{x}$$

$$8. \frac{d(a^x)}{dx} = a^x \log a$$



Question

Find the derivative of the following functions:

1. $f(x) = x^3 + 9x^2 - 48x + 2$

2. $f(x) = y^{-4} - 9y^{-3} + 8y^{-2} + 12$

Solution

$$\begin{aligned} 1. \quad f'(x) &= 3x^2 + 18x - 48 \\ &= 3(x^2 + 6x - 16) \\ &= 3(x+8)(x-2) \end{aligned}$$

$$\begin{aligned} 2. \quad f'(x) &= -4y^{-4-1} - 9(-3)y^{-3-1} + 8(-2)y^{-2-1} + 0 \\ &= -4y^{-5} + 27y^{-4} - 16y^{-3} \end{aligned}$$

2 Example

1. A toy company can sell x electronic gaming systems at a price of $p = -0.01x + 400$ dollars per gaming system. The cost of manufacturing x systems is given by $C(x) = 100x + 10,000$ dollars. Find the rate of change of profit when 10,000 games are produced. Should the toy company increase or decrease production?

Solution :

The profit $P(x)$ earned by producing x gaming systems is $R(x) - C(x)$,

Where,

$R(x)$ is the revenue obtained from the sale of x games. Since the company can sell x games at $p = -0.01x + 400$ per game,

$$\begin{aligned} R(x) &= x p \\ &= x(-0.01x + 400) \\ &= -0.01 x^2 + 400x. \end{aligned}$$

Consequently,

$$P(x) = -0.01 x^2 + 300x - 10,000.$$

2 Example

Therefore, evaluating the rate of change of profit gives :

$$\begin{aligned}
 P'(10000) &= \lim_{x \rightarrow 10000} \frac{P(x) - P(10000)}{x - 10000} \\
 &= \lim_{x \rightarrow 10000} \frac{-0.01x^2 + 300x - 10000 - 1990000}{x - 10000} \\
 &= \lim_{x \rightarrow 10000} \frac{-0.01x^2 + 300x - 2000000}{x - 10000} \\
 &= 100.
 \end{aligned}$$

Since the rate of change of profit $P'(10,000) > 0$ and $P(10,000) > 0$, the company should increase production.

3 Higher-Order Derivatives

The second derivative of a function is just the derivative of its first derivative. The third derivative is the derivative of the second derivative, the fourth derivative is the derivative of the third, and so on. The derivative of $f'(x)$, which is generally referred to as the second derivative of $f(x)$ and written $f''(x)$ or $f^{(2)}(x)$.

For example,

The derivative of a position function is the rate of change of position, or velocity. The derivative of velocity is the rate of change of velocity, which is acceleration.

3 Example

1. $f(x) = x^3$

Solution:

First derivative is

$$f'(x) = 3x^2$$

Second derivative is

$$f''(x) = 6x$$



Question

Find the second derivative of the following function:

1. $y = (4x^2 + 1)^2.$

Solution

$$y = (4x^2 + 1)(4x^2 + 1) = 16x^4 + 8x^2 + 1$$

First expand the brackets.

$$\frac{dy}{dx} = 64x^3 + 16x$$

Differentiate each of the three terms.

$$\frac{d^2y}{dx^2} = 192x^2 + 16$$

Then differentiate again.

4 Differentiation rules

1. Sum rule:

$$f(x) = g(x) + h(x)$$

Then,

$$f'(x) = g'(x) + h'(x).$$

4 Example

1. $f(x) = x + x^3$

Solution:

$$f'(x) = u'(x) + v'(x)$$

Now, differentiating the given function, we get;

$$f'(x) = \frac{d}{dx} (x + x^3)$$

$$f'(x) = \frac{d}{dx} (x) + \frac{d}{dx} (x^3)$$

$$f'(x) = 1 + 3x^2$$

4 Differentiation rules

2.Product Rule:

$f(x) = u(x) \times v(x)$,
then:



$$f'(x) = u'(x) \times v(x) + u(x) \times v'(x)$$

4 Example

1. $f(x) = x^2(x+3)$

Solution:

$$f'(x) = u'(x) \times v(x) + u(x) \times v'(x)$$

Here,

$$u(x) = x^2 \text{ and } v(x) = x+3$$

Therefore, on differentiating the given function, we get;

$$f'(x) = \frac{d}{dx} [x^2(x+3)]$$

$$f'(x) = \frac{d}{dx} (x^2)(x+3) + x^2 \frac{d}{dx} (x+3)$$

$$f'(x) = 2x(x+3) + x^2(1)$$

$$f'(x) = 2x^2 + 6x + x^2$$

$$f'(x) = 3x^2 + 6x$$

$$f'(x) = 3x(x+2)$$

4 Differentiation rules

3. Quotient Rule:

$$f(x) = u(x)/v(x)$$



$$\frac{u'(x) \times v(x) - u(x) \times v'(x)}{(v(x))^2}$$

4 Example

1. $\therefore k(x) = \frac{5x^2}{4x+3}.$

Solution:

$$k'(x) = \frac{f'(x)g(x) - g'(x)f(x)}{(g(x))^2} = \frac{10x(4x+3) - 4(5x^2)}{(4x+3)^2}.$$

$$k'(x) = \frac{20x^2 + 30x}{(4x+3)^2}.$$

5 Chain Rule of Differentiation

If a function $y = f(x) = g(u)$ and if $u = h(x)$, then the chain rule for differentiation is defined as;


$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

5 Example

1. $f(x) = (6x^2 + 7x)^4$

Solution:

$$\begin{aligned} f'(x) &= 4(6x^2 + 7x)^3 (12x + 7) \\ &= 4(12x + 7)(6x^2 + 7x)^3 \end{aligned}$$



Question

Solve the following function and find its first derivative:

1. $y = \sqrt{4 - 3x}.$

Solution

Let $u = 4 - 3x$

$$\frac{du}{dx} = -3$$

$$y = \sqrt{u} = u^{\frac{1}{2}}$$

$$\frac{dy}{du} = \frac{1}{2} u^{-\frac{1}{2}}$$

$$= \frac{1}{2} (4 - 3x)^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} = \frac{1}{2} (4 - 3x)^{-\frac{1}{2}} \times -3$$

$$= -\frac{3}{2} (4 - 3x)^{-\frac{1}{2}}$$

$$= -\frac{3}{2\sqrt{4 - 3x}}$$

Let u = the expression in the square root.

Now substitute $u = 4 - 3x$.

Apply the chain rule.

Simplify your answer.

6 Partial Differentiation

The process of finding the partial derivative of a function is called partial differentiation. In this process, the partial derivative of a function with respect to one variable is found by keeping the other variable constant. A partial derivative is defined as a derivative in which some variables are kept constant and the derivative of a function with respect to the other variable can be determined.

- The partial derivative of a function "f" with respect to "x" is represented by f_x or $\partial f / \partial x$.
- The partial derivative of a function "f" with respect to "y" is represented by f_y or $\partial f / \partial y$.

6 Example

Determine the partial derivative of the function: $f(x,y) = 3x + 4y$.

Solution:

Given function: $f(x,y) = 3x + 4y$

To find $\partial f / \partial x$, keep y as constant and differentiate the function:

Therefore, $\partial f / \partial x = 3$

Similarly, to find $\partial f / \partial y$, keep x as constant and differentiate the function:

Therefore, $\partial f / \partial y = 4$



Question

Find the derivative of the following function with respect to x and y:

1. $f(x, y) = x^4 y^3 + 8x^2 y + y^4 + 5x$

Solution

$$\frac{\partial z}{\partial x} = 4x^3y^3 + 16xy + 5 \quad (\text{Note: } y \text{ fixed, } x \text{ independent variable, } z \text{ dependent variable})$$

$$\frac{\partial z}{\partial y} = 3x^4y^2 + 8x^2 + 4y^3 \quad (\text{Note: } x \text{ fixed, } y \text{ independent variable, } z \text{ dependent variable})$$