



Subject : Statistical & Risk Modelling -2

Time Inhomogeneous Markov Jump

Agenda

- ✓ 1. Time In-homogenous Markov jump process
- ✓ 2. Chapman Kolmogorov equation
- ✓ 3. Kolmogorov Forward differential equations
- ✓ 4. Kolmogorov ~~Forward~~ ^{Backward} differential equations
- ✓ 5. Occupancy probability
- ✓ 6. Probability of going to another state
- ✓ 7. Integrated form of KFD and KBD
- ✓ 8. Applications

$$\mu = 0.5$$

$$\mu = 0.9 + 0.1k$$

Res: $d-1$ Holding Time

Time Inhomogeneity

$$\frac{25-30}{\text{age}} \rightarrow P_{HS}(25, 30)$$

$$P_{HS}(5, 6)$$

Markov jump process

Aged 20

min. stay age

In this chapter we discuss time-inhomogeneous Markov jump processes. The transition probabilities $P(X_t = j | X_s = i)$ for a time-inhomogeneous process depend not only on the length of the time interval $[s, t]$, but also on the times s and t when it starts and ends. This is because the transition rates for a time-inhomogeneous process vary over time.

pt. in time
from

$$\lambda(s) \quad \mu_{ij}(s)$$

trans. intensity
from Healthy
to Dead.

$$HSD \rightarrow P_{HS}(5) \rightarrow \text{Time homogeneous.}$$

$$P_{HS}(20, 25) \neq P_{HS}(50, 55)$$

Chapman Kolmogorov Eqns

The more general continuous-time Markov jump process $\{X_t, t \geq 0\}$ has transition probabilities:

$$p_{ij}(s, t) = P[X_t = j | X_s = i] \quad (s \leq t)$$

which obey a version of the *Chapman-Kolmogorov equations*, written in matrix form as:

$$P(s, t) = P(s, u)P(u, t) \quad \text{for all } s < u < t$$

or equivalently:

$$p_{ij}(s, t) = \sum_{k \in S} p_{ik}(s, u) p_{kj}(u, t) \quad \text{for all } s < u < t$$

$$p_{ij}(t) = \sum_k p_{ik}^{(s)} p_{kj}(t-s)$$

$$p_{ij}(s, t) = \sum_{k \in S} p_{ik}(s, u) \cdot p_{kj}(u, t)$$

$$s < u < t$$

Kolmogorov Eqns updated

Kolmogorov's forward differential equations (time-inhomogeneous case)

Written in matrix form these are:

$$\frac{\partial}{\partial t} P(s, t) = P(s, t) A(t)$$

where $A(t)$ is the matrix with entries $\mu_{ij}(t)$.

Kolmogorov's backward differential equations (time-inhomogeneous case)

The matrix form of Kolmogorov's backward equations is:

$$\frac{\partial}{\partial s} P(s, t) = -A(s)P(s, t)$$

Prove using Chapman Kolmogorov equation

Kolmogorov Eqns updated


μ γ r

Question

Write down Kolmogorov's forward and backward differential equation for $p_{HD}(s,t)$ & $p_{HS}(s,t)$.

KFD:

$$\begin{aligned} \text{KFD:} \\ \frac{dP_{HD}(s,t)}{dt} &= P_{HH}(s,t) \cdot \mu(t) \\ &+ P_{HS}(s,t) \cdot r(t) \end{aligned}$$



$$\frac{dP_{HD}(s,t)}{dt} = -\sigma(s) \cdot P_{HD}(s,t) + [\mu(s) + \sigma(s)] P_{HD}(s,t)$$

$$\frac{dP_{HS}(s,t)}{dt} = -\gamma(s) \cdot P_{HS}(s,t) + [\mu(s) + \sigma(s)] P_{HS}(s,t)$$

$$\frac{dP_{HS}(s,t)}{dt} = P_{HH}(s,t) \cdot \sigma(t) - P_{HS}(s,t) [\rho(t) + \gamma(t)]$$

Other probabilities

$$\lambda_i(u) = (0.01 + 0.005u)$$

$$H \rightarrow H - 6m$$

$$P_{ii}(s, t) = \exp\left(-\int_s^t \lambda_i(u) du\right)$$

$X_s < X_t = i$

Occupancy probabilities for time-inhomogeneous Markov jump processes

For a time-inhomogeneous Markov jump process:

$$p_{ii}(s, t) = \exp\left(-\int_s^t \lambda_i(u) du\right) = \exp\left(-\int_0^{t-s} \lambda_i(s+u) du\right)$$

where $\lambda_i(u)$ denotes the total force of transition out of state i at time u .


$$\lambda_i = \sum_j \mu_{ij}$$

$$P_{ii}(s, t) \rightarrow X_s = i, X_t = i$$

$$P_{ii}(s, t) \rightarrow X_s = i, X_t = i$$

Prove using Chapman Kolmogorov equation

Care of continuously staying in same state



Time-homogeneous
 $v_i = 0.9$

$$e^{-\lambda v_i}$$

Survival prob.

$$0.5 \int_0^{\infty} e^{-\lambda v_i} dv_i$$

Other probabilities

Question

A Markov jump process is used to model sickness and death. Four states are included, namely H, S_1, S_2 and D , which represent healthy, sick, terminally sick and dead, respectively. We are told that the people who are terminally sick never recover and die at a rate of $1.03(1.01)^t$ where t is their age in years.

Calculate the probability that a terminally sick 50-year-old dies within a year.

$$\begin{aligned}
 P_{S_2 D}(50, 51) &= 1 - P_{S_2 S_2}(50, 51) \Rightarrow P_{S_2 S_2}(50, 51) \\
 &= 1 - P_{S_2 S_2}(50, 51) = 1 - \exp\left(-\int_{50}^{51} 1.03(1.01)^t dt\right)
 \end{aligned}$$

$$n^t = \frac{n^{t+1}}{t+1}$$

e^{-N}
↓
decreases
↓
trans out
of state

$P_{S_2 D}$


Other probabilities

$$a^n = \frac{a^n}{\ln a}$$

Question

$$\begin{aligned}
 &= 1 - \exp \left[-1.03 \left(\frac{1.01^{51} - 1.01^{50}}{\ln 1.01} \right) \right] \\
 &= 1 - \exp \left[-1.03 \left(\frac{1.01^a}{\ln 1.01} \right) \right] \\
 &= 1 - \exp \left[-1.03 \left(\frac{1.01^{51} - 1.01^{50}}{\ln 1.01} \right) \right] \\
 &= \boxed{0.81725}
 \end{aligned}$$

Other probabilities



Probability that the process goes into state j when it leaves state i

Given that the process is in state i at time s and it stays there until time $s+w$, the probability that it moves into state j when it leaves state i at time $s+w$ is:

$$\frac{\mu_{ij}(s+w)}{\lambda_i(s+w)} = \frac{\text{the force of transition from state } i \text{ to state } j \text{ at time } s+w}{\text{the total force out of state } i \text{ at time } s+w}$$

$$\frac{p_{ij}(s+w)}{\lambda_i(s+w)}$$

$s \rightarrow$ starting time
 $w \rightarrow$ waiting time

$$\sum_{i \neq j} p_{ij} = \lambda_i$$



Residual Holding time

$$p_{ii}(s, t) = \exp\left(-\int_s^t \lambda_i(u) du\right) = \exp\left(-\int_0^{t-s} \lambda_i(s+u) du\right)$$

For a general Markov jump process, $\{X_t, t \geq 0\}$, define the residual holding time R_s as the (random) amount of time between s and the next jump:

$$\{R_s > w, X_s = i\} = \{X_u = i, s \leq u \leq s+w\}$$

$$P[R_s > w | X_s = i] = \exp\left(-\int_s^{s+w} \lambda_i(t) dt\right)$$



R_s is residual holding time
 R_s is staying in same state
 $P_{ii}(s, s+R_s) =$

Residual Holding time

$$\lambda(t) \rightarrow 0$$

$$\lambda_{12} = \mu_{12} + \mu_{12} \theta$$

pdf of R_s :

$$P(R_s > \omega) = 1 - \underbrace{P(R_s \leq \omega)}_{\text{cdf of } R_s}$$

$$\text{cdf of } R_s = 1 - \exp \left[- \int_s^{\infty} \lambda(u) \cdot du \right]$$

$$\text{pdf of } R_s : \frac{d F_R(\omega)}{d \omega} = - \frac{d \left(\exp \left[- \int_s^{\infty} \lambda(u) \cdot du \right] \right)}{d \omega}$$

$$\therefore \int \lambda(u) = \Delta(u)$$

$$\boxed{\int \lambda(u) = \Delta(u)}$$

Residual Holding time

$$\begin{aligned}
 &= - \frac{d \left[\exp \left[-(\Lambda(s+\omega) - \Lambda(s)) \right] \right]}{d\omega} \quad \text{cap. 1.2.1} \\
 &= - \exp \left[-(\Lambda(s+\omega) - \Lambda(s)) \right] \cdot \frac{d \left(\exp \left[-\Lambda(s+\omega) \right] \right)}{d\omega} \\
 &= - \exp \left[-(\Lambda(s+\omega) - \Lambda(s)) \right] \cdot \left(-\Lambda(s+\omega) \right) \\
 &\rightarrow \underline{\underline{\Lambda(s+\omega) \cdot \exp \left[-\int_s^{s+\omega} \lambda(u) \cdot du \right]}} \quad \leftarrow \text{sum}
 \end{aligned}$$

pdf of holding time

$$\lambda e^{-\lambda u}$$

Current Holding time

for how long are you in the current state?

For a full justification of this equation one needs to appeal to the properties of the current holding time C_t , namely the time between the last jump and t :

$$\{C_t \geq w, X_t = j\} = \{X_u = j, t - w \leq u \leq t\}$$

$$\downarrow$$

$$(C_t \geq w, X_t = j)$$

(t is your latest
R)

Integrated form of Kolmogorov Backward Eqn




$$p_{ij}(s, t) = \sum_{l \neq i} \int_0^{t-s} e^{-\int_s^{s+w} \lambda_i(u) du} \mu_{il}(s+w) p_{lj}(s+w, t) dw$$

provided $j \neq i$.

$$p_{ij}(s, t) = \sum_{k \neq i} \int_0^{t-s} p_{ii}(s, s+w) \cdot \mu_{ik}(s+w) \cdot p_{kj}(s+w, t) dw$$

$$p_{ii}(s, s+w) = \frac{p_{ii}(s, t)}{p_{ii}(s, t)}$$

- the probability of remaining in state i from time s to time $s+w$
- then making a transition to state k at time $s+w$
- and finally going from state k at time $s+w$ to state j at time t .



Integrated form of Kolmogorov Forward Eqn

$$p_{ij}(s, t) = \sum_{k \neq j} \int_0^{t-s} p_{ik}(s, t-w) \mu_{kj}(t-w) p_{jj}(t-w, t) dw$$


for $j \neq i$.

The factors in the integral are:


- the probability of going from state i at time s to state k at time $t-w$
- then making a transition from state k to state j at time $t-w$
- and staying in state j from time $t-w$ to time t .

$\int_{s_0}^s \cdot d\tau$
age

$\int_0^{t-s} \cdot d\tau$
duration



$\int_0^{t-s} p_{ik}(s, t-w) \cdot \mu_{kj}(t-w) \cdot p_{jj}(t-w, t) \cdot dw$



Exam style question

- ✓ (i) State the condition needed for a Markov Jump Process to be time inhomogeneous. [1]
- ✓ (ii) Describe the principal difficulties in modelling using a Markov Jump Process with time inhomogeneous rates. [2]

A multi-tasking worker at a children's nursery observes whether children are being 'Good' or 'Naughty' at all times. Her observations suggest that the probability of a child moving between the two states varies with the time, t , since the child arrived at the nursery in the morning. She estimates that the transition rates are:

From Good to Naughty: $0.2 + 0.04t$

From Naughty to Good: $0.4 - 0.04t$

where t is measured in hours from the time the child arrived in the morning, $0 \leq t \leq 8$.

trans. rates of
MJ process vary
with time.

Estimating rates
is more difficult
→ Solving diff. eqn
is comparatively
difficult

→ Data required
for modelling are
higher.

Exam style question

A child is in the 'Good' state when he arrives at the nursery at 9am.

- (iii) Calculate the probability that the child is Good for all the time up until time t . [3]
- (iv) Calculate the time by which there is at least a 50% chance of the child having been Naughty at some point. [2]

Let $P_G(t)$ be the probability that the child is Good at time t .

- (v) Derive a differential equation just involving $P_G(t)$ which could be used to determine the probability that the child is Good on leaving the nursery at 5pm. [2]


→ 13 mins

→ 3 mins

→ $P_G(t)$

$\frac{dP_G(t)}{dt}$



3 mins



Sick trans. order
→ d^2 of 't' and '(r)' → current holding time

$$s - (s - \omega) = \omega$$

$$\frac{u - (s - \omega)}{u + \omega - s} = \frac{\omega}{s}$$



$$t - (s - w)$$

$$= t + w - s$$

$$P[X_t = S, R_s > t-s \mid X_s = S, C_s = w] = \exp \left[- \int_s^t (\rho(u, w-s+u) + \nu(u, w-s+u)) du \right]$$

$$55 \times 2 = 110$$

$$= \exp \left[-\frac{1}{s} \right] \rho(u, \omega - s + u)$$

time spent
in sick state


$P \rightarrow S \rightarrow U$
 $V \rightarrow S \rightarrow D$

Sickness and Death with Duration

As a final example, the probability of being healthy at time t given that you are sick at time s with current illness duration w can be written as:

$$p_{S_w H}(s, t) = P[X_t = H \mid X_s = S, C_s = w]$$

$$= \int_s^t e^{-\int_s^v (\rho(u, w-s+u) + \nu(u, w-s+u)) du} \rho(v, w-s+v) p_{HH}(v, t) dv$$



$p_{S_{HH}}(s, t)$

$$= \int_s^t \exp \left[-\int_s^v (\rho(u, w-s+u) + \nu(u, w-s+u)) du \right] \rho(v, w-s+v) p_{HH}(v, t) dv$$

$$\cdot \rho(v, w-s+v)$$

$$\cdot p_{HH}(v, t) dv$$

$$\int p_{SS}(s, u) \cdot \rho(v, w-s+u) \cdot p_{HH}(v, t) du$$

Sickness and Death with Duration

Consider again the marriage model above, only now assume that the transition rate $d(t)$ depends on the current holding time. (So the chance of divorce depends on how long a person has been married.) Write down expressions for the probability that:

- (i) a bachelor remains a bachelor throughout a period $[s, t]$
- (ii) a person who gets married at time $s - w$ and remains married throughout $[s - w, s]$, continues to be married throughout $[s, t]$
- (iii) a person is married at time t and has been so for at least time w , given that they were divorced at time $s < t - w$.

Thank You