

Q1]

- i) A stochastic model is one that recognises the random nature of the input components. Stochastic models have the following advantages over deterministic models:
1. To reflect reality as accurately as possible, the model should imitate the random nature of the variables involved.
 2. A stochastic model can provide information about the distribution of the results (e.g. probabilities, variances etc), not just a single best estimate figure.
 3. Stochastic models allow you to use Monte Carlo simulation, which is an extremely powerful technique for solving complex problems.
- ii) For a time-inhomogeneous model the transition rates are functions of t . It is certainly possible to improve the fit by using a time-inhomogeneous model in this instance. However, If the age profile is represented by a density function $f(a)$; then the overall average rate at which a healthy employee falls sick is $= \lambda$, roughly constant for all t . The same of course applies to the overall average rate of recovery.
- iii) Proof: Given that Increment $X_{t+u} - X_t$ for every $u > 0$ are independent of past values of X_m and nonoverlapping.

Therefore, $Y = P[X_t = a \mid X_{S_1} = x_1, X_{S_2} = x_2, \dots, X_{S_n} = x_n, X_S = x]$

Because $X_S = x \rightarrow X_t = X_t - X_S + x$ --- (i)

$Y = P[X_t - X_S + x = a \mid X_{S_1} = x_1, X_{S_2} = x_2, \dots, X_{S_n} = x_n, X_S = x]$

Given $[X_t - X_S]$ are independent of past values of X , therefore $X_t - X_S + x$ will also be independent of past values of X_m .

Therefore:

$Y = P[X_t - X_S + x = a \mid X_S = x]$

$Y = P[X_t = a \mid X_S = x];$ Because $X_S = x$

Therefore Independent Increments satisfies Markov property

Q2]


- i) Consider a Markov chain taking values in the set $S = \{i : i = 0, 1, 2, 3, 4\}$, where i represents the number of umbrellas in the place where the Actuary currently is (at home or office).

If $i = 1$ and it rains then he take the umbrella, move to the other place, where there are already 3 umbrellas, and, including the one he brings, therefore he will now have 4 umbrellas. Thus, $p_{1,4} = p$, because p is the probability of rain.

If $i = 1$ but does not rain then he do not take the umbrella, goes to the other place and find 3 umbrellas. Thus, $p_{1,3} = 1 - p \equiv q$.

However, if $i=0$, he must move to other place where 4 umbrellas are kept with probability $P_{0,4}=1$

Similarly for other states. The process is depicted by the following diagram.



- ii) The transition matrix is as follow:

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & q & p \\ 0 & 0 & q & p & 0 \\ 0 & q & p & 0 & 0 \\ q & p & 0 & 0 & 0 \end{bmatrix}$$

- iii) For stationary distribution $\pi P = \pi$

Writing the equations:

$$\begin{aligned} \pi(0) &= \pi(4).q & \text{----1} \\ \pi(1) &= \pi(3).q + \pi(4).p & \text{----2} \\ \pi(2) &= \pi(2).q + \pi(3).p & \text{----3} \\ \pi(3) &= \pi(1).q + \pi(2).p & \text{----4} \\ \pi(4) &= \pi(0) + \pi(1).p & \text{----5} \end{aligned}$$

Substituting 1 in 5

$$\begin{aligned} \pi(4) &= \pi(4).q + \pi(1).p \Rightarrow \pi(1) = \pi(4). (1-q)/p \\ \Rightarrow \pi(1) &= \pi(4). p/p \Rightarrow \pi(1) = \pi(4) & \text{--- 6} \end{aligned}$$

Substituting 6 in 2

$$\pi(1) = \pi(3).q + \pi(4).p \Rightarrow \pi(4) = \pi(3).q + \pi(4).p \\ \Rightarrow \pi(3) = \pi(4).(1-p)/q \Rightarrow \pi(3) = \pi(4) \quad \text{---7}$$

Substituting 7 in 3

$$\pi(2) = \pi(2).q + \pi(3).p \Rightarrow \pi(2) = \pi(3).p/(1-q) \\ \Rightarrow \pi(2) = \pi(3).p/(1-q) \Rightarrow \pi(2) = \pi(3). \quad \text{---8}$$

Substituting 6 & 7 in 4

$$\pi(3) = \pi(1).q + \pi(2).p \Rightarrow \pi(4) = \pi(1).q + \pi(4).p \\ \Rightarrow \pi(1) = \pi(4).(1-p)/q \Rightarrow \pi(1) = \pi(4)$$

Therefore $\pi(1) = \pi(2) = \pi(3) = \pi(4) = \pi(0)/q$

But $\pi(0) + \pi(1) + \pi(2) + \pi(3) + \pi(4) = 1$, substituting in terms of $\pi(4)$
 $\pi(4).q + 4\pi(4) = 1$

Therefore $\pi(4) = \frac{1}{4+q} = \pi(1) = \pi(2) = \pi(3)$ and $\pi(0) = \frac{q}{4+q}$

- iv) He will wet every time he happens to be in state 0 and it rains. The chance he is in state 0 is $\pi(0)$. The chance it rains is p . Hence Probability that he gets wet.

$P(\text{WET})$ = Probability of being in state 0 and its raining

$$P(\text{WET}) = \frac{q}{4+q}.p$$

- v) $P(\text{WET}) = 0.6 \cdot 0.4 / (4 + 0.4) = 5.45\%$

If he want the chance to be less than 1% then, clearly, he need more umbrellas. So, suppose he need N umbrellas. Set up the Markov chain as above and generalising, it is clear that

$$\pi(1) = \pi(2) = \pi(3) = \pi(4) \dots \dots = \pi(N) = \pi(0)/q.$$

But $\pi(0) + \pi(1) + \pi(2) + \pi(3) + \pi(4) \dots \pi(N) = 1$, substituting in terms of $\pi(N)$

$$\Rightarrow \pi(N).q + N\pi(N) = 1$$

$$\Rightarrow \pi(N) = \frac{1}{N+q}$$

Therefore probability of getting Wet $P(\text{WET}) = p \cdot \pi(0) = \frac{p \cdot q}{N+q}$

$$\text{For } P(\text{WET}) < 1/100 \Rightarrow \frac{p \cdot q}{N+q} < 1/100$$

$$N > 100 \cdot p \cdot q - q ; \text{ given } p = 0.6, q = 0.4$$

$$N > 24 - 0.4 = 23.6$$

Thus it is not worth to keep 24 umbrellas instead of 4 to reduce the probability of getting wet from 6% to 1%. On the days he did not take the umbrella but it starts raining he may buy a cheap (use and throw) umbrella from nearby local shop or take a cab/ other mode of transport to office or borrow/ share an umbrella from some friend/ colleague using the same route or wait for the rain to subside etc. (any other suitable alternative is fine).

Q3]

i), ii)

	State Space	Time Set
Counting Process	Discrete	Discrete or Continuous
General Random Walk	Discrete or Continuous	Discrete
Poisson Process	Discrete	Continuous
Markov Jump Chain	Discrete	Discrete
Markov Jump Process	Discrete	Continuous

iii) a) Number of times the account has been overdrawn since it was opened – Poisson Process / Counting Process

b) Status (overdrawn, in credit) of the account on the last day of each month – Markov Jump Chain/ Counting Process

c) number of direct debits paid since the account was opened – Poisson


d) Status (overdrawn, in credit) of the account at any time since the account was opened. Markov Jump Process.

Q4]

i) Past history is needed to decide where to go in the chain. If a sportsman is at A and his/her performance reduces, you need to know what level of performance he was at the previous year to determine whether he or she drops one or two levels.

ii) The B level needs to be split into two. B+ is the level with no reduction in performance parameter last year B- is the level with reduction in performance parameter last year These levels are for modelling purposes and not the actual levels.

iii) The Process diagram is given below.



iv) The transition matrix is given by:

	D	C	B+	B-	A
D	0.5	0.5	0	0	0
C	0.3	0.2	0.5	0	0
B+	0	0.3	0.2	0	0.5
B-	0.3	0	0.2	0	0.5
A	0	0	0	0.3	0.7

iv) Stationary distribution:

The stationary distribution is the set of probabilities that satisfy the matrix equation

$\pi = \pi P$ with and additional condition $\sum \pi \pi_{ii} = 1$

Written out fully, this set of matrix equation is

$$0.5\pi\pi_{DD} + 0.3\pi\pi_{CC} + 0.3\pi\pi_{BB-} = \pi\pi_{DD} \text{ ---a)}$$

$$0.5\pi\pi_{DD} + 0.2\pi\pi_{CC} + 0.3\pi\pi_{BB+} = \pi\pi_{CC} \text{ ---b)}$$

$$0.5\pi\pi_{CC} + 0.2\pi\pi_{BB+} + 0.2\pi\pi_{BB-} = \pi\pi_{BB+} \text{ ---c)}$$

$$0.3\pi\pi_{AA} = \pi\pi_{BB-} \text{ ---d)}$$

$$0.5\pi\pi_{BB+} + 0.5\pi\pi_{BB-} + 0.7\pi\pi_{AA} = \pi\pi_{AA} \text{ ---e)}$$

$$\text{Substituting (d) in (e) } 0.5\pi\pi_{BB+} + 0.5 * 0.3\pi\pi_{AA} + 0.7\pi\pi_{AA} = \pi\pi_{AA} \Rightarrow \pi\pi_{BB+} =$$

$$0.3\pi\pi_{AA} \text{(vi)}$$

Substituting $\pi\pi_{BB+}$ aaaaaaa $\pi\pi_{AA}$ in (c) we get

$$0.5\pi\pi_{CC} + 0.2 * 0.3\pi\pi_{AA} + 0.2 * 0.3\pi\pi_{AA} = 0.3\pi\pi_{AA} \Rightarrow \pi\pi_{CC} = 0.36\pi\pi_{AA}$$

Substituting $\pi\pi_{BB+}$, $\pi\pi_{CC}$ aaaaaaa $\pi\pi_{AA}$ in (b)

$$0.5\pi\pi_{DD} + 0.2 * 0.36\pi\pi_{AA} + 0.3 * 0.3\pi\pi_{AA} = 0.36\pi\pi_{AA} \text{ ---b)}$$

$$\Rightarrow \pi\pi_{DD} = 0.396\pi\pi_{AA}$$

Substituting all in $\sum \pi \pi_{ii} = 1$

$$\pi\pi_{AA} + 0.3\pi\pi_{AA} + 0.3\pi\pi_{AA} + 0.36\pi\pi_{AA} + 0.396\pi\pi_{AA} = 1$$

$$\square \pi\pi_{AA} = 0.42445$$

$$\square \pi\pi_{BB-} = 0.12733$$

$$\square \pi\pi_{BB+} = 0.12733$$

$$\square \pi\pi_{CC} = 0.15280$$

$$\square \pi\pi_{DD} = 0.16808$$

vi) Long run average contract value is $100\% \pi_D + 120\% \pi_C + 150\% (\pi_{B+} + \pi_{B-}) + 175\% \pi_A = 1.4762$ million USD

v) Let m_i be the number of transitions (and years) taken to reach level A from any current level i. Then

$$m_{DD} = 1 + 0.5m_{DD} + 0.5m_{CC} \text{(a)}$$

$$m_{CC} = 1 + 0.2m_{CC} + 0.5m_{BB+} + 0.3m_{DD} \text{(b)}$$

$$m_{BB+} = 1 + 0.2m_{BB+} + 0.5m_{AA} + 0.3m_{CC} \text{(c)}$$

$$mmBB- = 1 + 0.3mmDD + 0.5mmAA + 0.2mmBB+(c)$$

$$mmAA = 0$$

from (a) we get,

$$mmDD = 2 + mmCC$$

substituting in (b) we get,

$$mmcc = 1 + 0.2mmCC + 0.5mmBB+ + 0.3(2 + mmCC)$$

$$mmBB+ = mmCC - 3.2$$

substituting in (c) ,

$$mmBB+ = 1 + 0.2mmBB+ + 0.5mmAA + 0.3mmCC$$

$$mmCC - 3.2 = 1 + 0.2(mmCC - 3.2) + 0.3mmCC$$

$$mmCC = 7.12 \text{ years}$$

Therefore, $mmDD = 2 + mmCC = 9.12 \text{ years}$

Q5]

- i) *There is an explicit dependence on the past behaviour of $(Y_j : j \leq n)$ in the probability distribution of Y_{n+1} ; further, X_n is nothing but the sum of Y_j .
Hence the markov property does not hold*
- ii) *In part i) we show that there is explicit dependence on the past behaviour of $(Y_j : j \leq n)$ and hence the markov property does not hold, this implies that the sequence $(Y_j : n \geq 1)$ does not form a markov chain.*
- iii) *The transition matrix is given by:*

$$\begin{vmatrix} p & 1-p & 0 & . & . & 0 \\ 0 & pe^{-\lambda} & 1-pe^{-\lambda} & 0 & . & . \\ . & 0 & pe^{-\lambda} & 1-pe^{-\lambda} & 0 & . \\ . & . & . & . & . & . \\ . & . & . & . & . & . \\ 0 & . & . & . & . & . \end{vmatrix}$$

- iv)
- The chain is time homogeneous since the transition probabilities calculated in part i) is independent of time n .*
 - It is irreducible, since the no. of errors can never go down.*
 - There are no recurrent states; hence there can be no stationary distribution. Alternatively, if a stationary distribution π exists, it has to follow:*

$$\begin{aligned} \pi_0 p &= \pi_0 \\ \pi_0(1-p) + \pi_1 pe^{-\lambda} &= \pi_2 \\ \pi_1(1-pe^{-\lambda}) + \pi_2 pe^{-2\lambda} &= \pi_2 \\ &\text{and so on.} \end{aligned}$$

*since $p \leq 1$, we have $\pi_0 = 0$, and then $\pi_1 = 0$,
Hence, no stationary distribution exists.*

- v) *Probability of no further error is*

$$(pe^{-j\lambda})^n = p^n e^{-nj\lambda}$$

Q6]

a). A stochastic model allows for the randomness of the input parameters.

Stochastic model have following advantage over deterministic model:

- a stochastic model provides the distribution of the results (probabilities and variances) and not just a single best estimate.
- Stochastic model correctly reflects the random nature of the variables involved as against deterministic one.
- Stochastic model allow to use Monte Carlo simulation which is a powerful technique to solve complex problem.

b). (i) Assume that the functions $p_{ij}(s, t)$ are continuously differentiable, the transition rates are defined by differentiation with respect to t .

$$\sigma_{ij}(s) = \left[\frac{d}{dt} (p_{ij}(s, t)) \right]_{t=s}$$

(ii) Since

$$\sum_{j \in S} p_{ij}(s, t) = 1 \text{ where } S - \text{State Space}$$

$$\begin{aligned} \text{We have } \sum_{j \in S} \sigma_{ij}(s) &= \sum_{j \in S} \left[\frac{d}{dt} p_{ij}(s, t) \right]_{t=s} \\ &= \frac{d}{dt} \sum_{j \in S} p_{ij}(s, t) \\ &= \frac{d}{dt} (1) \\ &= 0 \end{aligned}$$

Q7]

(a) The process may be expressed as a Markov chain by considering following states. Each state indicates the current number of successive defeats.

State Number of successive defeats

1 0 – Not defeated last time

2 1 – Defeated in the last match

3 2 – Defeated in the last two matches

4 3 – Defeated in the last three matches

5 4 – Captain sacked

The transition matrix is as follows:

$$\begin{bmatrix} 0.7 & 0.3 & 0 & 0 & 0 \\ 0.7 & 0 & 0.8 & 0 & 0 \\ 0.7 & 0 & 0 & 0.8 & 0 \\ 0.7 & 0 & 0 & 0 & 0.8 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

(b) A Markov chain is said to be irreducible if any state j can be reached from any state i .

The above process is not irreducible as the captain, once sacked, can never become the captain again.

(c)

i) The probability of remaining the captain for exactly four matches is given by:

Match	1	2	3	4	Probability
Result	Lose	Lose	Lose	Lose	
Probability	0.3	0.3	0.3	0.3	$0.3^4 = 0.0081$

ii) The probability of remaining captain for exactly 5 matches is given by

Match	1	2	3	4	5	Probability
Result	Lose	Lose	Lose	Lose	Lose	
Probability	0.3	0.3	0.3	0.3	0.3	$0.7 \times 0.3^4 = 0.00567$

iii) The probability of remaining captain for exactly 7 matches is given by

Match	1	2	3	4	5	6	7	Probability
Result	Any	Any	Not Lose	Lose	Lose	Lose	Lose	
Probability	1	1	0.3	0.3	0.3	0.3	0.3	$0.7 \times 0.3^4 = 0.00567$

iv) The probability of remaining captain for exactly 9 matches is given by

Match	1	2	3	4	5	6	7	8	9	Probability
Result	Not all four losses				Not Lose	Lose	Lose	Lose	Lose	
Probability	$= 1 - 0.3^4 = 0.9919$				0.7	0.3	0.3	0.3	0.3	$= 0.005624$

(d)

Define N_1

= Expected number of matches as a captain given that the current state is i .
Since, the captain is newly appointed, the variable N in the question is N_1 as defined here.

we have:

$$N_1 = 1 + 0.7 * N_1 + 0.3 * N_2$$

$$N_2 = 1 + 0.7 * N_1 + 0.3 * N_3$$

$$N_3 = 1 + 0.7 * N_1 + 0.3 * N_4$$

$$N_4 = 1 + 0.7 * N_1$$

Solving the above equations ,we get;

$$N_4 = 123.46$$

$$N_3 = 160.49$$

$$N_2 = 171.60$$

$$N_1 = 174.94$$

$$\text{therefore, } E[N] = 174.94$$

Q8]

- (i) The score currently stands at 'Tie'. Whosever wins the next point will move into a 'Lead'. If the player in 'Lead' wins the subsequent point as well, he would win the tie-breaker. However, if the player in 'Lead' loses the next point, the score would be back at 'Tie'.

Since the probability of moving to the next state does not depend on the history prior to entering the state, Markov property holds.

The state space is defined as follows:

State	Description
T	Tie
L _F	Federer Leads
L _N	Nadal Leads
G _F	Federer Wins
G _N	Nadal Wins

- (ii) The transition matrix is set out below:

$$\begin{bmatrix} 0 & 0.55 & 0.45 & 0 & 0 \\ 0.45 & 0 & 0 & 0.55 & 0 \\ 0.55 & 0 & 0 & 0 & 0.45 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

- (iii) The chain is reducible as it has two absorbing states – G_F and G_N.

Absorbing states have no period and the other three states have a period of 2. Thus, the chain is not a-periodic.

- (iv) After two points from the tie, the tie-breaker would either be completed or be back to tie again.

The probability of returning to tie after two points is given by:

Probability of Federer winning the first point x Probability of Nadal winning the second point
+
Probability of Nadal winning the first point x Probability of Federer winning the second point

$$= 0.55 \times 0.45 + 0.45 \times 0.55 \\ = 0.495$$

We need to find number of such cycles of returning to tie such that

$$0.495^N = 1 - 0.95$$

Solving the above equation:

$$N = \frac{\ln 0.05}{\ln 0.495} = 4.26$$

Since the game can finish in cycles of two points, the required number of cycles is 5 i.e. 10 points.

- (v) After two points:
- Nadal may have won the tie-breaker (probability of 0.2025 i.e. 0.45^2); or
 - Federer may have won the tie-breaker (probability of 0.3025 i.e. 0.55^2); or
 - Tie-breaker may have come back to tie (probability of 0.495).

Let F_T be the probability that Federer wins the tie-breaker that is presently tied.

Let N_T be the probability that Nadal wins the tie-breaker that is presently tied.

We have:

$$N_T = 0.2025 + 0.495 \times N_T$$

Solving:

$$N_T = 0.401$$

Probability that Federer eventually wins the tie-breaker is 0.599 ($1 - N_T$). This can be verified by:

$$F_T = 0.3025 + 0.495 \times F_T$$


Solving:

$$F_T = 0.599$$

Probability of Nadal winning a point is 0.45. However, in order to win the game, Nadal would need to win at least two consecutive points at some point in the game. The probability of Nadal winning two consecutive points is lower than the probability of him winning a point – this is what one would reasonably expect.

Q9]

a) The state transition diagram is set out below:



b) We are to calculate for the % of corporate buyer having a target % for XYZ of 65%

- In 2 years time
- Over the long-run.

The state of the system after one year $S_1 = S_0 P$

$$(0.05, 0.30, 0.45, 0.20) \begin{pmatrix} 0.6 & 0.3 & 0.1 & - \\ - & 0.7 & 0.3 & - \\ - & 0.4 & 0.4 & 0.2 \\ - & 0.2 & 0.5 & 0.3 \end{pmatrix} = (0.03, 0.445, 0.375, 0.15)$$

Hence the state of the system in 2 years time $S_2 = S_1 P$

$$(0.03, 0.445, 0.375, 0.15) \begin{pmatrix} 0.6 & 0.3 & 0.1 & - \\ - & 0.7 & 0.3 & - \\ - & 0.4 & 0.4 & 0.2 \\ - & 0.2 & 0.5 & 0.3 \end{pmatrix} = (0.018, 0.5005, 0.3615, 0.12)$$

Hence the % of corporate buyer having a target % of 65% for XYZ in 2 years time is 36.15%.

The long run steady state can be found by solving the following equation:

$$S = SP$$

$$(x_1, x_2, x_3, x_4) = (x_1, x_2, x_3, x_4)P$$

$$(x_1, x_2, x_3, x_4) = (x_1, x_2, x_3, x_4) \begin{pmatrix} 0.6 & 0.3 & 0.1 & - \\ - & 0.7 & 0.3 & - \\ - & 0.4 & 0.4 & 0.2 \\ - & 0.2 & 0.5 & 0.3 \end{pmatrix}$$

Hence we have the five equations

$$x_1 = 0.6x_1$$

$$x_2 = 0.3x_1 + 0.7x_2 + 0.4x_3 + 0.2x_4$$

$$x_3 = 0.1x_1 + 0.3x_2 + 0.4x_3 + 0.5x_4$$

$$x_4 = 0.2x_3 + 0.3x_4$$

$$x_1 + x_2 + x_3 + x_4 = 1$$

Now from the first equation above,

$$0.4x_1 = 0$$

So

$$x_1 = 0$$

Substituting this into the other equations above and rearranging we get

$$0.3x_2 = 0.4x_3 + 0.2x_4 \quad (1)$$

$$0.6x_3 = 0.3x_2 + 0.5x_4 \quad (2)$$

$$0.7x_4 = 0.2x_3 \quad (3)$$

$$x_2 + x_3 + x_4 = 1 \quad (4)$$

One of the above equations is redundant.

Hence we have

$$(x_1 = 0, x_2 = 0.5423, x_3 = 0.3559, x_4 = 0.1017)$$

- c)** The steady state proportion of customers having a 65% target allocation for XYZ Ltd is 35.59%.

It is noted that the initially this proportion is 45% and it quickly drops to:

- 37.5% in one year's time; and
- 36.2% in two years' time

It is, therefore, likely that the steady state shall be reached in a few years' time.

Q10]

- i. If $i = 1$ and it's *raining*, then the shopkeeper takes his umbrella and moves to the other place. When he reaches the other place (shop or home as the case may be), he will end up with three umbrellas at that place.

If p is the probability of *raining* then the shopkeeper will carry the umbrella with probability p if there are 1 or 2 umbrellas from where the shopkeeper is presently starting.

If there are three umbrellas from where the shopkeeper is presently starting, then the probability of carrying umbrella is $p' = 0.25(1 + 3p)$ and probability of not carrying an umbrella is $0.75(1 - p)$, then.

$P_{i,j}$ is the probability of carrying umbrella from state i to state j .

Thus,

$$P_{1,3} = p$$

If $i = 1$ and it's not a *raining* and shopkeeper is not carrying an umbrella, he goes to the other place and find 2 umbrellas. Thus,

$$P_{1,2} = q$$

where $q = 1 - p$

And starting from 3 umbrellas,

$$P_{3,1} = p' = 0.25(1 + 3p)$$

$$P_{3,0} = q' = 0.75(1 - p)$$

Continuing in the same manner I form a Markov chain with the following diagram:



The transition matrix is given by :

$$\begin{matrix} & 0 & 1 & 2 & 3 \\ \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & q & p \\ 0 & q & p & 0 \\ q' & p' & 0 & 0 \end{pmatrix} \end{matrix}$$

ii. In order to find stationary distribution we solve the system of equations

$$\begin{aligned} \pi P &= \pi \\ \pi_0 + \pi_1 + \pi_2 + \pi_3 &= 1 \end{aligned}$$

$$\pi_0 = q' \pi_3$$

$$\pi_3 = \pi_0 + p\pi_1$$

$$\pi_1 = q \pi_2 + p' \pi_3$$

$$\pi_2 = q \pi_1 + p\pi_2$$

Solving the last equation,

We get $\pi_1 = \pi_2$

Also,

$$\pi_1 = \frac{.25(1+3p)\pi_3}{p} = \pi_2$$

Using this in the $\sum \pi$ we get,

$$2\pi_1 + \pi_3 + q'\pi_3 = 1$$

$$2 \frac{.25(1+3p)\pi_3}{p} + \pi_3 + 0.75q\pi_3 = 1$$

$$\pi_3 = \frac{p}{2 \times .25(1+3p) + p + 0.75pq}$$

$$\pi_0 = \frac{0.75(1-p)}{2 \times .25(1+3p) + p + 0.75pq}$$

Every time the shopkeeper gets wet when he happens to be in state 0 and it's raining. The probability that he is in state 0 is π_0 and the probability that it's raining, is p .

Hence, the required probability is

$$p \cdot \pi_0 = \frac{0.75(1-p)p}{2 \times .25(1+3p) + p + 0.75pq}$$

iii.

Now we know that, $p = 0.7$

Hence, the required probability is 4.58%

If I want the chance to be less than 2% then, clearly, I need more umbrellas.

So, suppose he needs N umbrellas. By following the same Markov chain (as above) for N umbrellas, we find that

$$\pi_{N-1} = \pi_{N-2} \dots \dots \dots = \pi_2 = \pi_1$$

Also

$$\pi_1 = \frac{0.25(1+3p)\pi_N}{p} = \pi_2 \dots \dots = \pi_{N-1}$$

And

$$\pi_0 = q' \pi_N$$

Using this in the $\sum \pi$ we get,

$$(N-1) \pi_1 + \pi_N + q' \pi_N = 1$$

$$(N-1) \frac{0.25(1+3p)}{p} \pi_N + \pi_N + q' \pi_N = 1$$

$$(N-1)0.25(1+3p) \pi_N + p\pi_N + pq' \pi_N = p$$

$$\pi_N = \frac{p}{(N-1) \times .25(1+3p) + p + 0.75pq}$$

Hence,

$$\pi_0 = \frac{0.75p(1-p)}{(N-1) \times .25(1+3p) + p + 0.75pq}$$

The Probability that the shopkeeper gets wet $\pi_0 p$, which is

$$\frac{0.75(1-p)p^2}{(N-1) \times .25(1+3p) + p + 0.75pq}$$

As we want the required probability to be less 2%,

$$0.75(1-p)p^2 < 2\% \{ (N-1) \times .25(1+3p) + p + 0.75pq \}$$

$$75(1-p)p^2 < 2 \{ (N-1) \times .25(1+3p) + p + 0.75pq \}$$

$$75(1-p)p^2 < 0.5(N-1)(1+3p) + 2p + 1.5pq$$

$$75(1-p)p^2 - 2p - 1.5pq < 0.5(N-1)(1+3p)$$

$$1 + \frac{75(1-p)p^2 - 2p - 1.5pq}{0.5(1+3p)} < N$$

Solving the above equation gives $N=7$.

So to reduce the chance of getting wet to less than 2% the shopkeeper needs 7 umbrellas.

Q11]

i) If each room is represented by the state, then the transition matrix P for this Markov chain is as follows:

$$P = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1/4 & 1/4 & 0 & 1/4 & 1/4 & 0 \\ 0 & 0 & 1/2 & 0 & 0 & 1/2 \\ 0 & 0 & 1/2 & 0 & 0 & 1/2 \\ 0 & 0 & 0 & 1/2 & 1/2 & 0 \end{pmatrix}$$

ii) The chain is irreducible, because it is possible to go from any state to any other state. However, it is not aperiodic, because for any even n , $P^n_{6,1}$ will be zero and for any odd n $P^n_{6,5}$ will also be zero. This means that there is no power of P that would have all its entries strictly positive.

iii) For P to be stationary,

$$\pi P = P$$

Perform matrix multiplication and show that πP is equal to P

iv) We find from π that the mean recurrence time (i.e. the expected time to return) for the room 1 is

$$1/\pi(1)=12$$

v) Let, $\psi(i) = E(\text{number of steps to reach state 5} \mid X_0 = i)$.

We have

$$\psi(5) = 0$$

$$\psi(6) = 1 + (1/2)\psi(5) + (1/2)\psi(4)$$

$$\psi(4) = 1 + (1/2)\psi(6) + (1/2)\psi(3)$$

$$\psi(3) = 1 + (1/4)\psi(1) + (1/4)\psi(2) + (1/4)\psi(4) + (1/4)\psi(5)$$

$$\psi(1) = 1 + \psi(3)$$

$$\psi(2) = 1 + \psi(3).$$

We solve and find $\psi(1) = 7$.

Q12]

i)

We wish to find m_{24}

$$\text{Now, } m_{i4} = \begin{cases} 0, & \text{if } i = 4, \\ 1 + \sum_{j \neq 4} p_{ij} m_{j4}, & \text{if } i \neq 4 \end{cases}$$

$$\text{Thus, } m_{24} = 1 + (1 \times m_{34})$$

$$m_{44} = 0$$

$$m_{34} = 1 + \left(\frac{1}{3} \times m_{44}\right) + \left(\frac{2}{3} \times m_{24}\right)$$

$$m_{34} = 1 + 0 + \frac{2}{3} (1 + m_{34})$$

$$m_{34} = 5$$

$$\text{Hence, } m_{24} = 1 + 5 = 6 \text{ steps}$$

ii)

$$P(1,2,3,2,3,4) = P(X_0 = 1) \times (p_{12}) \times (p_{23}) \times (p_{32}) \times (p_{23}) \times (p_{34})$$

$$= \frac{3}{4} \times \frac{3}{5} \times 1 \times \frac{2}{3} \times 1 \times \frac{1}{3}$$

$$= \frac{1}{10}$$

Q13]

i)

a) Data : RSRSSSSRRSRSSRRSRSSSSRRSSRRR

$$p_{rr} = 6/14 = 3/7$$

$$p_{rs} = 8/14 = 4/7$$

$$p_{sr} = 8/15$$

$$p_{ss} = 7/15$$

b)

30 th June	1 st Jul	2 nd Jul	3 rd Jul	
R	S	S	R	
	$p_{rs} = 8/14$	$p_{ss} = 7/15$	$p_{sr} = 8/15$	0.142222
R	S	R	R	
	$p_{rs} = 8/14$	$p_{sr} = 8/15$	$p_{rr} = 6/14$	0.130612
R	R	S	R	
	$p_{rr} = 6/14$	$p_{rs} = 8/14$	$p_{sr} = 8/15$	0.130612
R	R	R	R	
	$p_{rr} = 6/14$	$p_{rr} = 6/14$	$p_{rr} = 6/14$	0.078717

Total = 0.482164

ii)

a)

Here, $\mu = 3$

"Some policies " means "1 or more policies" i.e 1 minus the "zero policies" probability:

$$P(X > 0) = 1 - P(x_0)$$

$$\text{Now, } P(X) = \frac{e^{-\mu} \mu^x}{x!}$$

$$\text{So, } P(x_0) = \frac{e^{-3} 3^0}{0!} = 4.9787 \times 10^{-2}$$

Therefore the probability of 1 or more policies is given by:

$$\begin{aligned} \text{Probability} &= P(X \geq 0) \\ &= 1 - P(x_0) \\ &= 1 - 4.9787 \times 10^{-2} \\ &= 0.95021 \end{aligned}$$

b)

The probability of selling 2 or more, but less than 5 policies is:

$$P(2 \leq X < 5) = P(x_2) + P(x_3) + P(x_4)$$

$$= \frac{e^{-3} 3^2}{2!} + \frac{e^{-3} 3^3}{3!} + \frac{e^{-3} 3^4}{4!}$$

$$= 0.61611$$

c)

Average number of policies sold per day: $\frac{3}{5} = 0.6$

$$\text{So on a given day, } P(X) = \frac{e^{-0.6} 0.6^1}{1!} = 0.32929$$