

Numerical Method & Algebra

Assignment - 1

Q1

$$\begin{aligned}\text{Actual length} &= x_1 = 12.5 \text{ cm} \\ \text{Actual area} &= \pi r_1^2 \\ &= \pi (12.5)^2 \\ &= 490.87365 \text{ cm}^2\end{aligned}$$

$$\begin{aligned}x_2 &= 12.65 \text{ cm} \\ \text{Approximate area} &= \pi r_2^2 \\ &= \pi (12.65)^2 \\ &= 502.72551 \text{ cm}^2\end{aligned}$$

$$\begin{aligned}\text{Absolute error} &= \text{App. area} - \text{Actual area} \\ &= 11.84886\end{aligned}$$

$$\begin{aligned}\text{Relative error} &= \frac{\text{Absolute area}}{\text{Actual Area}} \\ &= 0.02413\end{aligned}$$

$$\begin{aligned}\text{Percentage error} &= (\text{Relative error} \times 100) \\ &= 2.418 \%\end{aligned}$$

Q2 i)

$$\begin{aligned}x_1 &= 4 & y_1 &= 0.60206 \\ x_2 &= 6 & y_2 &= 0.7781513\end{aligned}$$

Using Linear Interpolation

$$x_2 - x_1 = y_2 - y_1$$

$$\begin{aligned}(6 - 4) &= 0.7781913 - 0.60206 \\ 2 &= 0.1760913\end{aligned}$$

$$\text{Charge} = \frac{0.1760913}{2}$$

$$= 0.8804565$$

Thus, $n = 5$

$$y = 0.60206 + 0.8804565$$

$$y = 0.69010565$$

ii) $x_1 = 4.5$

$$y_1 = 0.6532125$$

$$x_2 = 5.5$$

$$y_2 = 0.7403627$$

$$(5.5 - 4.5) = y_2 - y_1$$

$$= 0.7403627 - 0.6532125$$

$$= 0.0871502$$

$$\text{Charge} = \frac{0.0871502}{1}$$

Thus $n = 5$

$$y = 0.6532125 + 0.0871502$$

$$= 0.6967876$$

Q3. $x^4 - x - 10 = 0$ $a = 1.5, b = 2$

x	y
$a = x_0 = 1.5$	-6.4375
$b = x_1 = 2$	4

Numerical Method & Algebra

Assignment - 1

Q1

$$\begin{aligned} \text{Actual length} &= x_1 = 12.5 \text{ cm} \\ \text{Actual area} &= \pi r_1^2 \\ &= \pi (12.5)^2 \\ &= 490.87365 \text{ cm}^2 \end{aligned}$$

$$\begin{aligned} x_2 &= 12.65 \text{ cm} \\ \text{Approximate area} &= \pi r^2 \\ &= \pi (12.65)^2 \\ &= 502.72551 \text{ cm}^2 \end{aligned}$$

$$\begin{aligned} \text{Absolute error} &= \text{App. area} - \text{Actual area} \\ &= 11.84886 \end{aligned}$$

$$\begin{aligned} \text{Relative error} &= \frac{\text{Absolute area}}{\text{Actual Area}} \\ &= 0.02413 \end{aligned}$$

$$\begin{aligned} \text{Percentage error} &= (\text{Relative error} \times 100) \\ &= 2.418 \%. \end{aligned}$$

Q2 i)

$$\begin{aligned} x_1 &= 4 & y_1 &= 0.60206 \\ x_2 &= 6 & y_2 &= 0.7781513 \end{aligned}$$

Using Linear Interpolation

$$x_2 - x_1 = y_2 - y_1$$

$$\begin{aligned} (6 - 4) &= 0.7781913 - 0.60206 \\ 2 &= 0.1760913 \end{aligned}$$

$$\text{Change} = \frac{0.1760913}{2}$$

$$= 0.8804565$$

Thus, $n = 5$

$$y = 0.60206 + 0.8804565$$

$$y = 0.69010565$$

ii)

$$x_1 = 4.5$$

$$y_1 = 0.6532125$$

$$x_2 = 5.5$$

$$y_2 = 0.7403627$$

$$(5.5 - 4.5) = y_2 - y_1$$

$$= 0.7403627 - 0.6532125$$

$$= 0.0871502$$

$$\text{Change} = \frac{0.0871502}{1}$$

Thus $n = 5$

$$y = 0.6532125 + 0.0871502$$

$$= 0.6967876$$

Q3. $x^4 - x - 10 = 0$ $a = 1.5, b = 2$

x	y
$a = x_0 = 1.5$	-6.4375
$b = x_1 = 2$	4

$$1. \quad n_2 = \frac{n_0 + n_1}{2} = 1.75$$

$$y_2 = -2.37109$$

$$2. \quad n_3 = \frac{n_1 + n_2}{2}$$

$$= 1.875$$

$$y_3 = 0.48462$$

$$3. \quad n_4 = \frac{n_2 + n_3}{2}$$

$$= 1.8125$$

$$y_4 = -1.0202$$

$$4. \quad n_5 = \frac{n_4 + n_3}{2}$$

$$= 1.84375$$

$$y_5 = -0.28773$$

$$5. \quad n_6 = \frac{n_5 + n_3}{2}$$

$$= 1.859375$$

$$y_6 = 0.0953$$

$$Q4. \quad n^3 - n - 1 = 0$$

$$f'(n) = 3n^2 - 1$$

n	y
0	1
1	-1
2	5

$$x_0 = 1$$

$$f(x_0) = -1$$

$$f'(x_0) = 2$$

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$= 1 - \left(\frac{-1}{2} \right)$$

$$= 1.5$$

$$y_1 = 0.875$$

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$$= 1.5 - \left(\frac{0.875}{0.75} \right)$$

$$x_2 = 1.34783$$

$$f(x_2) = 0.100699$$

$$f'(x_2) = 4.44998$$

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)}$$

$$= 1.3252$$

$$y_3 = 0.00209$$

$$f'(x_3) = 4.26846$$

$$x_4 = x_3 - \frac{f(x_3)}{f'(x_3)}$$

$$= 1.32472$$

$$y_4 = 0.000008712$$

Q5. $A^2 = A$

$$\begin{aligned} 7A - (I + A)^3 &= 7A - (I^3 + A^3 + 3A^2I + 3AI) \\ &= 7A - (I + A^2(A) + 3A + 3A) \\ &= 7A - (I + 7A) \\ &= -I \end{aligned}$$

Q6. $A = \begin{bmatrix} 1 & -1 & 2 \\ 4 & 0 & 6 \\ 0 & 1 & -1 \end{bmatrix}$

$$|A| = 0 - 1(6 - 8) + (-1)4$$

$$\text{Adj}(A) = \begin{bmatrix} -6 & 1 & -6 \\ 4 & -1 & 2 \\ 4 & -1 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj}(A)}{|A|}$$

$$= \frac{-1}{2} \begin{bmatrix} -6 & 1 & -6 \\ 4 & -1 & 2 \\ 4 & -1 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj}(A)}{|A|}$$

$$= \frac{-1}{2} \begin{bmatrix} -6 & 1 & -6 \\ 4 & -1 & 2 \\ 4 & -1 & 4 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} +3 & -1/2 & 3 \\ -2 & 1/2 & -1 \\ 2 & 1/2 & -2 \end{bmatrix}$$

Q 7

$$\begin{aligned} \vec{a} &= 8\hat{i} - 9\hat{j} \\ \vec{b} &= 7\hat{i} - 9\hat{j} \end{aligned}$$

$$\begin{aligned} \vec{a} \cdot \vec{b} &= ab \cos \theta \\ &= (8\hat{i} - 9\hat{j}) \cdot (7\hat{i} - 9\hat{j}) \\ &= 56 + 81 \\ &= 137 \end{aligned}$$

$$\begin{aligned} a &= \sqrt{64 + 81} = \sqrt{145} \\ b &= \sqrt{49 + 81} = \sqrt{130} \end{aligned}$$

$$\begin{aligned} \vec{a} \cdot \vec{b} &= ab \cos \theta \\ \cos \theta &= \frac{\vec{a} \cdot \vec{b}}{ab} \\ &= \frac{137}{\sqrt{130} \times \sqrt{145}} \end{aligned}$$

$$\cos \theta = 0.9720224206$$

Q 8.

$$\vec{R} = \vec{a} + \vec{b}$$

$$R^2 = A^2 + B^2 + 2AB \cos \theta$$

$$\theta = 30^\circ$$

$$R^2 = (75)^2 + (25)^2 + 2(75)(25) \cos 30^\circ$$

$$R^2 = 9497.595284$$

Q 9.

$$d = 3$$

$$AP = 101, 104, 107, \dots, 998$$

$$a = 101$$

$$a_n = a + (n-1)d$$

$$998 = 101 + (n-1)3$$

$$n = 300$$

$$S_n = \frac{n}{2} (a + e)$$

$$= \frac{300}{2} (101 + 998)$$

$$= 150 + 1099$$

$$S_n = 164850$$

Q 10.

$$a_6 = 32$$

$$a_n^5 = 32$$

$$\frac{a_n^7}{a_n^5} = \frac{128}{32}$$

$$r^2 = 4$$

$$r = \pm 2$$

$$r = 2$$

Q 13.

$$f(x) = x^3 - 7x^2 + 8x - 3$$

$$f'(x) = 3x^2 - 14x + 8$$

$$x = 5$$

$$f(5) = -13$$

$$f'(5) = 13$$

$$n_1 = n_0 - \frac{f(n_0)}{f'(n_0)}$$

$$= 5 - \left(\frac{-13}{13} \right)$$

$$= 6$$

$$f(x_1) = 9$$

$$f'(x_1) = 3x_1$$

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$$x_2 = 6 - 9/3 = 5$$

$$x_2 = 5.71875$$

$$f(x_2) = 0.84787$$

Q14. $(x^2 - x + 1)^4 = [(x-1)^2 + x]^4$

using Pascal's triangle

$$[(x-1)^2 + x]^4 = (x-1)^8 + 4(x-1)^6x + 6(x-1)^4x^2 + 4(x-1)^2x^3 + x^4$$

$$(x^2 - x + 1)^4 = (x-1)^8 + 4x(x-1)^6 + 6x^2(x-1)^4 + 4x^3(x-1)^2 + x^4$$

Q11. $(3x+4)^5 = (3x)^5 + 5(3x)^4 \cdot 4 + 10(3x^3) \cdot 4^2 + [10(3x^2) \cdot 4^3] + 5(3x) \cdot 4^4 + 4^5$

$$= 243x^5 + 1620x^4 + 4320x^3 + 5760x^2 + 3840x + 1024$$

Q12. $A = \begin{bmatrix} 2 & -3 & 0 \\ 2 & -5 & 0 \\ 0 & 0 & 3 \end{bmatrix}$

$$A - \lambda I = \begin{bmatrix} 2 & -3 & 0 \\ 2 & -5 & 0 \\ 0 & 0 & 3 \end{bmatrix} = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

$$= \begin{bmatrix} 2-\lambda & -3 & 0 \\ 2 & -5-\lambda & 0 \\ 0 & 0 & 3-\lambda \end{bmatrix}$$

$$|A - \lambda I| = -\lambda^3 + 13\lambda - 12$$

$$= -\lambda^3 + 13\lambda - 12 = 0$$

$$\lambda^3 - 13\lambda + 12 = 0$$

Comparing coefficient

$$\Rightarrow 1 - 13 + 12 = 0$$

$$\therefore \text{Root} = 1$$

$$\lambda^2 - \lambda - 12 = 0$$

$$\lambda^2 + 4\lambda - 3\lambda - 12 = 0$$

$$\lambda = 3 \quad \text{or} \quad -4$$

Eigen Value = -4, 1 & 3

Q15.

$$e^n = 3 \log(n)$$

$$3 \log(n) + e^{-n} = 0$$

x	y
0.1	-2.0951625
0.2	-1.278179
0.3	-0.8278180
0.4	-0.5235
0.5	-0.29656
0.6	-0.11673
0.7	-0.03188

$$x_0 = 0.6$$

$$y_0 = -0.11673$$

$$x_1 = 0.7$$

$$y_1 = 0.03188$$

$$x_2 = \frac{x_0 + x_1}{2} = 0.65$$

$$y_2 = -0.03921$$

$$x_3 = \frac{x_2 - x_1}{2} = 0.675$$

$$y_3 = -0.00293$$

$$x_4 = \frac{x_3 + x_1}{2}$$

$$= 0.6875$$

$$y_4 = 0.01465$$

$$x_5 = \frac{x_4 + x_3}{2} = 0.68125$$

$$y_5 = 0.0059$$