

1]

- (i) Writing the state space in the order {Bid (B), Offer (O)},
the generator matrix is:

$$\begin{matrix} B & \begin{pmatrix} -\lambda & \lambda \end{pmatrix} \\ O & \begin{pmatrix} \mu & -\mu \end{pmatrix} \end{matrix}.$$

- (ii) The holding times are exponentially distributed with
parameter λ in state B ,
and μ in state O .

(iii) $\frac{\partial}{\partial t} {}_tP_s^{BB} = -\lambda {}_tP_s^{BB} + \mu {}_tP_s^{BO}.$

$$\frac{\partial}{\partial t} {}_tP_s^{BO} = \lambda {}_tP_s^{BB} - \mu {}_tP_s^{BO}.$$

- (iv) We have a two-state model so:

$${}_tP_s^{BB} + {}_tP_s^{BO} = 1.$$

Substituting:

$$\frac{\partial}{\partial t} {}_tP_s^{BB} = -\lambda {}_tP_s^{BB} + \mu(1 - {}_tP_s^{BB});$$

$$\frac{\partial}{\partial t} \left[\exp((\lambda + \mu)t) \cdot {}_tP_s^{BB} \right] = \mu \cdot \exp((\lambda + \mu)t);$$

and hence


$$\exp((\lambda + \mu)t) \cdot {}_tP_s^{BB} = \frac{\mu}{\lambda + \mu} \cdot \exp((\lambda + \mu)t) + \text{constant}.$$

Since the process is in state Bid at time s (i.e. $t = 0$),
the constant is $\frac{\lambda}{\mu + \lambda}$,

$$\text{and thus } {}_tP_s^{BB} = \frac{\mu}{\lambda + \mu} + \frac{\lambda}{\lambda + \mu} \cdot \exp(-(\lambda + \mu)t).$$

2]

(i)



(ii)
$$\frac{d}{dt}P_{AA}(t) = -0.3P_{AA}(t) + 0.1P_{AB}(t) + 0.3P_{AC}(t)$$

$$\frac{d}{dt}P_{AB}(t) = 0.2P_{AA}(t) - 0.5P_{AB}(t) + 0.1P_{AC}(t)$$

$$\frac{d}{dt}P_{AC}(t) = 0.1P_{AA}(t) + 0.4P_{AB}(t) - 0.4P_{AC}(t)$$

(iii) EITHER

To stay in state A the equation reduces to:

$$\frac{d}{dt}P_{AA}(t) = -0.3P_{AA}(t)$$

$$P_{AA}(t) = \exp(-0.3t)$$

So for $t = 2$ we have $\exp(-0.6) = 0.5488$.

OR

We can model this as Poisson with parameter $(0.1 + 0.2) \cdot 2 = 0.6$

$$P(\text{Poi}(0.6) = 0) = \frac{e^{-0.6} 0.6^0}{0!}$$

$$= e^{-0.6} = 0.5488$$

- (iv) The only paths under which the third jump is into state C are BAC , CAC and CBC .

The probabilities of each jump are given by the ratio of the transition rates.

So the probabilities for each path are:

$$BAC = \frac{2}{3} \cdot \frac{1}{5} \cdot \frac{1}{3} = \frac{2}{45}$$


$$CAC = \frac{1}{3} \cdot \frac{3}{4} \cdot \frac{1}{3} = \frac{1}{12}$$

$$CBC = \frac{1}{3} \cdot \frac{1}{4} \cdot \frac{4}{5} = \frac{1}{15}$$

$$\text{Sum} = 7/36 = 0.194.$$

3]

(i)



(ii) Using the Markov assumption

OR

the Chapman Kolmogorov equation is

$${}_{dt+t}P_x^{34} = {}_tP_x^{31} {}_{dt}P_{x+t}^{14} + {}_tP_x^{32} {}_{dt}P_{x+t}^{24} + {}_tP_x^{33} {}_{dt}P_{x+t}^{34} + {}_tP_x^{34} {}_{dt}P_{x+t}^{44} + {}_tP_x^{35} {}_{dt}P_{x+t}^{54}.$$

Since ${}_{dt}P_{x+t}^{54} = {}_tP_x^{31} = 0$

$${}_{dt+t}P_x^{34} = {}_tP_x^{32} {}_{dt}P_{x+t}^{24} + {}_tP_x^{33} {}_{dt}P_{x+t}^{34} + {}_tP_x^{34} {}_{dt}P_{x+t}^{44}.$$

Given that ${}_{dt}P_{x+t}^{44} = 1$

And assuming that, for small dt

$${}_{dt}P_{x+t}^{ij} = \mu_{x+t}^{ij} dt + o(dt) \quad i \neq j$$

where $\lim_{dt \rightarrow 0} \frac{o(dt)}{dt} = 0$,

then substituting, we have

$${}_{dt+t}P_x^{34} = {}_tP_x^{32} \mu_{x+t}^{24} dt + {}_tP_x^{33} \mu_{x+t}^{34} dt + {}_tP_x^{34} + o(dt)$$

so that ${}_{dt+t}P_x^{34} - {}_tP_x^{34} = {}_tP_x^{32} \mu_{x+t}^{24} dt + {}_tP_x^{33} \mu_{x+t}^{34} dt + o(dt)$

and hence $\frac{d}{dt}({}_tP_x^{34}) = \lim_{dt \rightarrow 0} \frac{{}_{t+dt}P_x^{34} - {}_tP_x^{34}}{dt} = {}_tP_x^{32} \mu_{x+t}^{24} + {}_tP_x^{33} \mu_{x+t}^{34}.$

4]

- (i) The mean is equal to the parameter, so there are 3 calls per hour.
- (ii) The process is memoryless so the fact that Fred has not had a call for 15 minutes is irrelevant.

Expected time until next call is 20 minutes.

- (iii) This is the probability of zero calls in time 0.5 hours.

Using $p_j(t) = e^{-\lambda t} (\lambda t)^j / j!$

OR

$$\text{Since } p_0(0.5) = \frac{e^{-1.5} (1.5)^0}{0!},$$

$$p_0(0.5) = e^{-1.5} = 0.2231 .$$

- (iv) The expected time that Fred is on the phone is the expected number of calls times the expected length of a call.

Per hour this is 3 calls times 7 minutes = 21 minutes.

So the probability that the phone is engaged is $21/60 = 0.35$.

5]

(i) The Chapman Kolmogorov equation is

$$p_{HH}(x, t+dt) = p_{HH}(x, t)p_{HH}(t, t+dt) \\ + p_{HS}(x, t)p_{SH}(t, t+dt) + p_{HD}(x, t)p_{DH}(t, t+dt)$$

But $p_{DH}(t, t+dt) = 0$ or other explanation why path through D can be ignored

So:

$$p_{HH}(x, t+dt) = p_{HH}(x, t)p_{HH}(t, t+dt) + p_{HS}(x, t)p_{SH}(t, t+dt)$$

Assuming that, for small dt

$$p_{ij}(t, t+dt) = \lambda_{ij}(t)dt + o(dt) \quad i \neq j$$

$$p_{ii}(t, t+dt) = 1 + \lambda_{ii}(t)dt + o(dt)$$

OR

$$p_{ii}(t, t+dt) = 1 - \sum_{j \neq i} \lambda_{ij}(t)dt + o(dt)$$

where the λ s are the instantaneous transition rates and $\lim_{dt \rightarrow 0} \frac{o(dt)}{dt} = 0$,

then substituting, we have

$$p_{HH}(x, t+dt) = p_{HH}(x, t)(1 - \sigma(t)dt - \mu(t)dt) + p_{HS}(x, t)\rho(t, C_t) + o(dt)$$

so that

$$p_{HH}(x, t+dt) - p_{HH}(x, t) = p_{HH}(x, t)(-\sigma(t) - \mu(t))dt \\ + p_{HS}(x, t)\rho(t, C_t)dt + o(dt)$$

and hence

$$\frac{d}{dt} p_{HH}(x, t) = \lim_{dt \rightarrow 0} \frac{p_{HH}(x, t+dt) - p_{HH}(x, t)}{dt} \\ = p_{HH}(x, t)(-\sigma(t) - \mu(t)) + p_{HS}(x, t)\rho(t, C_t)$$

(ii) The equation simplifies when considering $p_{\overline{HH}}(t)$ to

$$\frac{d}{dt} p_{\overline{HH}}(0, t) = -(\sigma(t) + \mu(t)) p_{\overline{HH}}(t)$$

$$\frac{1}{p_{\overline{HH}}(0, t)} \frac{d}{dt} p_{\overline{HH}}(0, t) = -(\sigma(t) + \mu(t)) = \frac{d}{dt} \ln p_{\overline{HH}}(t) .$$

Integrate both sides:

$$\left[\ln p_{\overline{HH}}(0, t) \right]_0^t = \int_{s=0}^t -(\sigma(s) + \mu(s)) ds$$

as $p_{\overline{HH}}(0) = 1$

$$p_{\overline{HH}}(0, t) = \exp - \left(\int_{s=0}^t (\sigma(s) + \mu(s)) ds \right)$$

6]

All three processes have a discrete state space.

A Markov Chain and Markov Jump Chain both operate in discrete time but a Markov jump

Process operates in continuous time.

All have the Markov property which is EITHER that the future development of the process can be predicted from its present state alone, without reference to its past history.

$$P[X_t \in A \mid X_{s_1} = x_1, X_{s_2} = x_2, \dots, X_{s_n} = x_n, X_s = x] = P[X_t \in A \mid X_s = x]$$

for all times $s_1 < s_2 < \dots < s_n < s < t$, all states $x_1, x_2, \dots, x_n, x$ in S and all subsets A of S .

The Jump Chain obeys the Markov Property and behaves as a Markov Chain except when the Jump Chain encounters an absorbing state. From that time, it makes no further transitions, implying that time stops for the Jump Chain.

The Jump Chain associated with X takes the same path through the state space as X does.

The Markov Jump Chain and the Markov Chain are expressed in terms of probabilities whereas the Markov Jump Process is expressed in terms of rates.

The Markov Chain can have loops in each state, the Markov Jump process cannot and the Markov Jump Chain only has loops on absorbing states.

7]

- (i) The maximum likelihood estimate of the transition intensity from state i to state j is the number of transitions from state i to state j divided by the total waiting time in state i .

To estimate the transition intensities exactly we therefore need

the total time spent in each state

OR

entry and exit times for each individual for each state,

and the total number of transitions of each type made.

- (ii) Define $p_{AA}(s, t)$ to be the probability of being in state Active at time $s+t$ if Active at time s .

Then EITHER

$$\frac{\partial}{\partial t} p_{AA}(s, t) = -p_{AA}(s, t)\mu$$

$$\frac{\partial}{\partial t} p_{AT}(s, t) = p_{AA}(s, t)\mu ,$$

OR

$$\frac{\partial}{\partial t} p(s, t) = p(s, t)M$$

where $M = \begin{pmatrix} -\mu & \mu \\ 0 & 0 \end{pmatrix}$ in order Active, Theft,

OR

Integrated forward equations:

$$p_{AA}(s, t) = \exp\left(-\int_{u=s}^t \mu du\right)$$

$$p_{AT}(s, t) = \int_{u=0}^t p_{AA}(s, u) \mu \cdot 1 du .$$

- (iii) Measure from time zero i.e. $s = 0$ and drop s from notation.

EITHER

$$\frac{1}{p_{AA}(t)} \frac{\partial}{\partial t} p_{AA}(t) = -\mu .$$

$$\frac{\partial}{\partial t} (\ln(p_{AA}(t))) = -\mu ,$$

hence $p_{AA}(t) = \exp(-\mu t + C)$.

As $p_{AA}(0) = 1$, $C = 0$, so

$$p_{AA}(t) = \exp(-\mu t)$$

A claim occurs with cost $\pounds C$ if moves to state “Theft Claim”.

Hence the expected cost is $C(1 - \exp(-\mu T))$

OR

Solving for p_{AT} , we have

$$\frac{\partial}{\partial t} p_{AT}(t) = p_{AA}(t)\mu = (1 - p_{AT}(t))\mu \text{ (as the model has only two states).}$$

Using an integrating factor, we can write


$$\frac{\partial}{\partial t} [\exp(\mu t) p_{AT}(t)] = \mu \exp(\mu t) ,$$

$$\exp(\mu t) p_{AT}(t) = \exp(\mu t) - 1 ,$$

$$p_{AT}(t) = 1 - \exp(-\mu t) ,$$

and hence the expected cost is $C(1 - \exp(-\mu T))$.

(iv)



(v) We now have $\frac{\partial}{\partial t} p_{AA}(t) = -p_{AA}(t)(\mu + \lambda)$.

So $p_{AA}(t) = \exp(-(\mu + \lambda)t)$.

We want $\frac{\partial}{\partial t} p_{AT}(t) = p_{AA}(t)\mu = \mu \exp(-(\mu + \lambda)t)$.

Solving this produces


$$p_{AT}(t) = \left| \frac{-\mu}{(\mu + \lambda)} \exp(-(\mu + \lambda)t) \right|_0^T = \frac{\mu}{\mu + \lambda} (1 - \exp(-(\mu + \lambda)T)) .$$

So claims become $\frac{\mu}{\mu + \lambda} C(1 - \exp(-(\mu + \lambda)T))$.

8]

(i) $\{0,1,2,3,4,\dots\}$

(ii)



(iii) Generator matrix

Lives	0	1	2	3	4	...
	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & . \\ \mu & -(\mu+\lambda) & \lambda & 0 & 0 & \\ 0 & \mu & -(\mu+\lambda) & \lambda & 0 & \\ 0 & 0 & \mu & -(\mu+\lambda) & \lambda & \\ 0 & 0 & 0 & \mu & -(\mu+\lambda) & \\ . & . & . & . & . & . \end{pmatrix}$					

(iv) EITHER

If a Markov jump process X_t is examined only at the times of transition, the resulting process is called the jump chain associated with X_t .

OR

A jump chain is each distinct state visited in the order visited where the time set is the times when states are moved between.

(v)	Lives	0	1	2	3	4	...
		$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & \text{etc.} \\ \mu / (\mu + \lambda) & 0 & \lambda / (\mu + \lambda) & 0 & 0 & \\ 0 & \mu / (\mu + \lambda) & 0 & \lambda / (\mu + \lambda) & 0 & \\ 0 & 0 & \mu / (\mu + \lambda) & 0 & \lambda / (\mu + \lambda) & \\ 0 & 0 & 0 & \mu / (\mu + \lambda) & 0 & \\ \text{etc.} & & & & & \end{pmatrix}$					

(vi) $\left(\frac{\mu}{\mu + \lambda} \right)^3$

9]

(i)

- The transition rates of moving from each state to each other state must be non-negative
- **OR** The transition rates OFF the leading diagonal must be non-negative
OR $\mu_{ij} \geq 0$ for $i \neq j$
- The transition rates ON the leading diagonal must be non-positive
- **OR** $\mu_{ii} \leq 0$
- The sum of each row must be zero.
OR $\mu_{ii} = -\sum_{j \neq i} \mu_{ij}$
- Should be a finite or countable square matrix.

(ii) State space $\{2,1,0\}$ policies in force.

(iii)



(iv)

- $\frac{d}{dt} P_{22}(t) = -0.4 P_{22}(t)$
- $\frac{d}{dt} P_{21}(t) = 0.4 P_{22}(t) - 0.2 P_{21}(t)$
- $\frac{d}{dt} P_{20}(t) = 0.2 P_{21}(t)$

10]

$$(i) \quad \frac{d}{dt} P_{AA}(t) = -2t \times P_{AA}(t)$$

$$\Rightarrow \frac{d}{dt} [\ln P_{AA}(t)] = -2t$$

$$\Rightarrow \ln P_{AA}(s) = -s^2 + \text{constant}$$

We know $P_{AA}(0) = 1$, hence constant = 0

$$\text{Hence, } P_{AA}(s) = \exp^{-s^2}$$


$$(ii) \quad P(\text{in first visit to B at time } T \mid \text{in state A at } t = 0)$$

$$\begin{aligned} &= \int_0^T P(\text{remains in A to time } s) \\ &\quad \times P(\text{transition to B in time } s, s + ds) \\ &\quad \times P(\text{remains in B to time } T) ds \\ &= \int_{s=0}^T P_{AA}(s) \times 2s \times P_{BB}(s, T) ds \end{aligned}$$

Using the result from part (i) and the similar result for P_{BB} with boundary condition $P_{BB}(s, s) = 1$, this gives us:

$$\begin{aligned} &= \int_{s=0}^T e^{-s^2} \times 2s \times e^{-T^2+s^2} ds \\ &= \int_{s=0}^T 2s \times e^{-T^2} ds \\ &= e^{-T^2} \times T^2 \end{aligned}$$

(iii) (a) The sketch should be shaped like:



(b) Commentary:

- Initially probability increases from 0 at $T = 0$, and accelerates as the transition rate from A to B increases.
- However, as transitions increase, it becomes more likely that the process has already visited state B and jumped back to A. Therefore the probability of being in the first visit to B tends (exponentially) to zero.

(c) Differentiate to find turning point:

$$\frac{d}{dt} \left[e^{-t^2} \times t^2 \right] = 2t \times e^{-t^2} - 2t^3 \times e^{-t^2}$$

set derivative equal to zero

$$e^{-t^2} \times 2t \times (1 - t^2) = 0$$

implies $t = 1$ for a positive solution

and, from above analysis, this is clearly a maximum.

11]

- (i) Let N_t denote the number of claims up to time t . Since the Poisson process has stationary increments, we may take $t = 0$, so that the required conditional distribution is

$$\begin{aligned} P(T_0 \leq y | N_s = 1) &= \frac{P(T_0 \leq y, N_s = 1)}{P(N_s = 1)} \\ &= \frac{P(N_y = 1, N_s - N_y = 0)}{P(N_s = 1)} \end{aligned}$$

But $N_s - N_y$ is independent of N_y and has the same distribution as N_{s-y} .

Thus the right hand side above equals

$$\frac{(\lambda y e^{-\lambda y}) e^{-\lambda(s-y)}}{\lambda s e^{-\lambda s}} = \frac{y}{s},$$

which is the cdf of the uniform distribution on $[0, s]$.

- (ii) Since holding times are independent, each having an exponential distribution, their joint density is

$$\lambda^n e^{-\lambda(t_1 + t_2 + \dots + t_n)} 1_{\{t_1, t_2, \dots, t_n > 0\}}.$$

- (iii) We have, as in part (i),

$$\begin{aligned} P(N_s = k | N_t = n) &= \frac{P(N_s = k, N_t = n)}{P(N_t = n)} \\ &= \frac{P(N_s = k, N_t - N_s = n - k)}{P(N_t = n)} \end{aligned}$$

Using again that the Poisson process has stationary and independent increments, and that the number of claims in an interval $[0, t]$ is Poisson (λt) , we derive from above that

$$\begin{aligned}
 P(N_s = k \mid N_t = n) &= \frac{\frac{e^{-\lambda s} (\lambda s)^k}{k!} \cdot \frac{e^{-\lambda(t-s)} \lambda^{n-k} (t-s)^{n-k}}{(n-k)!}}{\frac{e^{-\lambda t} (\lambda t)^n}{n!}} \\
 &= \frac{e^{-\lambda t} \lambda^n s^k (t-s)^{n-k}}{k!(n-k)!} \cdot \frac{n!}{e^{-\lambda t} \lambda^n t^n} \\
 &= \frac{n!}{k!(n-k)!} \cdot \frac{s^k (t-s)^{n-k}}{t^k t^{n-k}} \\
 &= \binom{n}{k} \left(\frac{s}{t}\right)^k \left(1 - \frac{s}{t}\right)^{n-k}
 \end{aligned}$$

which is binomial with parameters n and s/t .